

2



INSTITUTE REPORT NO. 222

AD-A175 296

INTERACTION OF
PHENYLDICHLOROARSINE WITH BIOLOGICAL MOLECULES

RICHARD J. O'CONNOR, MS
EVELYN L. McGOWN, PhD
and
KILIAN DILL*, PhD

DIVISION OF COMPARATIVE MEDICINE AND TOXICOLOGY
LETTERMAN ARMY INSTITUTE OF RESEARCH

and

*DEPARTMENT OF CHEMISTRY
CLEMSON UNIVERSITY
CLEMSON, SC

DTIC FILE COPY

AUGUST 1986

LETTERMAN ARMY INSTITUTE OF RESEARCH
PRESIDIO OF SAN FRANCISCO, CALIFORNIA 94129

DTIC
ELECTE
DEC 22 1986
S E D

This document has been approved
for public release and sale; its
distribution is unlimited.

86 12 22 022

INTERACTION OF PHENYLDICHLOROARSINE WITH BIOLOGICAL MOLECULES


O'Connor, McGown, and Dill

Reproduction of this document in whole or in part is prohibited except with the permission of the Commander, Letterman Army Institute of Research, Presidio of San Francisco, California 94129. However, the Defense Technical Information Center is authorized to reproduce the document for United States Government purposes.

Destroy this report when it is no longer needed. Do not return it to the originator.

Citation of trade names in this report does not constitute an official endorsement or approval of the use of such items.

This material has been reviewed by Letterman Army Institute of Research and there is no objection to its presentation and/or publication. The opinions or assertions contained herein are the private views of the author(s) and are not to be construed as official or as reflecting the views of the Department of the Army or the Department of Defense. (AR 360-5)

 27 Aug 56
.....
(Signature and date)

This document has been approved for public release and sale; its distribution is unlimited.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER INSTITUTE REPORT NO. 222	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Interaction of Phenylldichloroarsine with Biological Molecules		5. TYPE OF REPORT & PERIOD COVERED Final - 1 May '85-1 May '86
7. AUTHOR(s) Richard J. O'Connor, MS; Evelyn L. McGown, PhD; and Kilian Dill, PhD		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Division of Comparative Medicine & Toxicology Letterman Army Institute of Research Presidio of San Francisco, CA 94129-6800		8. CONTRACT OR GRANT NUMBER(s)
11. CONTROLLING OFFICE NAME AND ADDRESS US Army Medical Research & Development Command Fort Detrick Frederick, MD 21701-5010		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Project #3M162734A875 ✓ Task BD Work Unit #306
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE August 1986
		13. NUMBER OF PAGES 19
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) This document is approved for public release and sale: distribution is unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Lewisite analogue; arsenical; phenylldichloroarsine; British anti-Lewisite; meso-2,3-dimercapto succinic acid; glutathione; ¹³ C-nuclear magnetic resonance.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) <u>Carbon 14</u> <u>Carbon 13</u> ¹⁴ C-Phenylldichloroarsine (PDA) was used to study the binding of PDA to erythrocytes and stroma-free hemoglobin. Natural abundance ¹³ C-nuclear magnetic resonance spectroscopy (¹³ C-NMR) was used to study the binding of PDA to glutathione, meso-2,3-dimercapto-succinic acid (DMSA), and British anti-Lewisite (BAL). The results clearly show that PDA penetrates the red		

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 65 IS OBSOLETE

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

cont'd

Cont 13
UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

cell and is contained within. Apparently hemoglobin is not the internal sulfhydryl-containing species that holds the PDA; glutathione appears to be a better candidate. ^{Carbon 13} NMR is a promising tool for investigating PDA complexes.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

ABSTRACT

¹⁴C-Phenyldichloroarsine (PDA) was used to study the binding of PDA to erythrocytes and stroma-free hemoglobin. Natural abundance ¹³C-nuclear magnetic resonance spectroscopy (¹³C-NMR) was used to study the binding of PDA to glutathione, meso-2,3-dimercaptosuccinic acid (DMSA), and British anti-Lewisite (BAL). The results clearly show that PDA penetrates the red cell and is contained within. Apparently hemoglobin is not the "internal" sulfhydryl-containing species that "holds" the PDA; glutathione appears to be a better candidate. ¹³C-NMR is a promising tool for investigating PDA complexes.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



PREFACE

Kilian Dill was a participant in the U.S. Army Summer Faculty Research and Engineering Program during summer of 1985. His current address is:

Department of Chemistry
Clemson University
Clemson, SC 29634-1905

ACKNOWLEDGMENTS

The authors would like to express their appreciation to CPT Mike McCreery for his help with the project.


TABLE OF CONTENTS

	Page
Abstract	i
Preface.	ii
Acknowledgments.	iii
Table of Contents.	v
BODY OF REPORT	
INTRODUCTION.	1
METHODS	2
PDA	2
PDA and Intact Red Blood Cells.	2
PDA and Human Hemoglobin.	3
Preparation of PDA Adducts.	3
NMR Studies	3
RESULTS AND DISCUSSION.	4
PDA Interaction with Intact Red Blood .	
Cells	4
PDA and Hemoglobin.	4
¹³ C-NMR Studies of PDA and its Adducts.	4
PDA	5
PDA and BAL	5
PDA and Glutathione	7
PDA and DMSA.	10
CONCLUSIONS	10
RECOMMENDATIONS	11
REFERENCES.	12
APPENDIX (Figures 1-4).	13
DISTRIBUTION	19

Interaction of Phenyldichloroarsine with Biological Molecules--O'Connor et al

INTRODUCTION

Phenyldichloroarsine (PDA) is a vesicant arsenical whose molecular toxicology is little understood (1). It is an analog of Lewisite (transchlorovinyl-dichloroarsine) and has a similar vesicant potency (2,3). In order to study the efficacy of various antidotes to these compounds, it is of paramount importance to study the molecular binding, transport, and the eventual cellular damage that PDA can produce.



Mono-substituted trivalent arsenicals such as PDA react with thiol groups to produce multiple monothiol species and with vicinal dithiol compounds to produce thermodynamically stable cyclic compounds (1). Because of the specific reactivity of PDA with thiol groups, it is likely that the biological damage caused by PDA results from the reaction with thiol-containing molecules present in biological systems.

In order to gain insight into the biological actions of PDA, with the goal of finding improved arsenic antidotes, we investigated the interaction of PDA with the red blood cell and its cytoplasmic thiol (SH)-containing molecules. Within the red cell, hemoglobin contributes the bulk of exposed free

SH groups, 1.8-3.6 $\mu\text{mol}/10^9$ cells, depending upon the species. The protein contains an additional 3.6-7.2 $\mu\text{mol SH}/10^9$ cells which are inaccessible unless the protein is denatured. Glutathione (GSH) contributes 0.15-0.30 $\mu\text{mol SH}/10^9$ cells (4). Red cell membranes and ergothioneine account for < 5% of the total free SH content (5).

In the following report we present our results with ^{14}C -PDA and mouse red blood cells and human hemoglobin. We also present nuclear magnetic resonance (NMR) spectroscopic analyses of PDA adducts with several small molecular weight SH compounds.

METHODS

PDA

Phenyldichloroarsine (PDA) was purchased from Research Organic/Inorganic Chemical Corp., Sun Valley, CA. It was purified by repeated distillation at 128°C under 15 mm Hg of nitrogen and the purity (> 99%) was verified by infrared spectrophotometry. Stock solutions of PDA were prepared at a concentration of 50 mM in anhydrous ethanol. ^{14}C -PDA (5.6 mCi/mmmole) was purchased from Wizard Laboratories, Davis, CA.

The aqueous form of PDA is easily oxidized to phenylarsonic acid. Our ^{14}C -PDA preparation was synthesized by a method which utilizes the oxidation of iodide ion to iodine to reduce the arsenical to its desired oxidation state (III). If trace amounts of iodine are left in the product, either free or as an adduct with PDA, it will promote oxidation of PDA when it is exposed to water. We found this was a significant problem with our commercial ^{14}C -PDA. To minimize the problem, we first distilled the ^{14}C -PDA under high vacuum, leaving behind the phenylarsonic acid (and any other non-volatile impurities). The PDA was then stored under vacuum to deprive it of oxygen and to maintain strictly anhydrous conditions. Stored in this manner, the isotope was stable for weeks.

PDA and Intact Red Blood Cells

In order to ascertain whether the red blood cell quantitatively takes up PDA and, if so, whether PDA

binds to the membrane or some cytoplasmic component, we undertook the radio-labelling of red blood cells with ^{14}C -PDA. Whole blood was obtained from six white mice (ICR strain; approximately 30 g) via heart puncture and centrifuged at $300 \times g$ for 30 minutes. The supernatant and buffy coat were removed, the cells resuspended and washed twice in saline. Packed cells were then resuspended with an equal volume of saline followed by the addition of the ^{14}C -PDA (approximately $0.25 \text{ } \mu\text{mol}/10^9$ cells). The reaction was allowed to proceed for two hours. The cells were then spun down, the supernatant saved, and the cells were lysed with distilled water. The membranes were pelleted by centrifugation and washed several times by resuspending the membranes in water, repelleting and removing the supernatant. The pellets and washings were also saved for counting.

PDA and Human Hemoglobin

To study PDA binding to human hemoglobin (Hb), stroma-free hemoglobin was obtained from the Blood Research Division, mixed with ^{14}C -PDA in varying proportions, put into dialysis bags and dialyzed against 40 ml saline. ^{14}C was determined in both the internal contents and external fluid at timed intervals.

Preparation of PDA Adducts

PDA-glutathione was prepared by the following method: PDA was added to a beaker containing 20 ml of anhydrous ethanol. To this a twofold molar excess of reduced glutathione was added in four equal portions over a period of approximately 15 minutes. Glutathione gradually dissolved in the ethanol as it reacted with PDA. The solution was continuously stirred for another hour and then evaporated to dryness using gaseous nitrogen. All other adducts of PDA were made by simply placing the appropriate molar amounts of PDA and DMSA or BAL into the desired solvent. Adducts were formed immediately.

NMR Studies

Proton-decoupled, natural abundance ^{13}C -NMR spectra were obtained on a Varian XL-300 FT-NMR spectrometer operating at 75.429 megahertz (7.05 tesla). A

25 KHz window and an 18 usec (90°) pulse width were used throughout these studies. The spectra of the PDA-glutathione adduct were obtained in a water/ deuterated water (3:1) mixture; PDA in D-4 methanol and D-6 acetone; the PDA-DMSA adduct in D-4 methanol and D-6 acetone; the PDA-BAL adduct in D-4 methanol and D-6 acetone.

RESULTS AND DISCUSSION

PDA Interaction with Intact Red Blood Cells

When we used freshly distilled ^{14}C -PDA, the intact red cells took up approximately 98% of the ^{14}C . When older ^{14}C preparations were used, the amount that remained extracellular increased, presumably due to the breakdown product, phenylarsonic acid. The latter compound was not taken up by red blood cells (R.J. O'Connor and E.L. McGown, unpublished results). When the cells were lysed, > 90% of the ^{14}C was in the cytoplasm and < 10% was membrane-associated.

PDA and Hemoglobin

The results above indicated that PDA penetrated the red cell membrane and was bound to some component in the cell cytoplasm. To investigate the possibility that hemoglobin was the cytoplasmic component to which PDA was bound, human hemoglobin-PDA equilibrium dialysis experiments were performed. It was found that 96% of the radiolabel had dialyzed out. We attempted to determine the number of "tight" binding sites per Hb molecule by using the Scatchard plot (Figure 1). There appeared to be only 0.05 sites per Hb molecule. These results indicated that PDA interacts weakly with Hb (0.05 sites) or that it does not bind to Hb (PDA may be binding to an impurity). Overall, we concluded that when PDA enters the red blood cell it binds to some component other than Hb, possibly glutathione.


^{13}C -NMR Studies of PDA and its Adducts

In the following discussion, chemical shifts are given relative to external tetramethylsilane (TMS); the respective solvents were used as internal references.

PDA. ^{13}C spectra of PDA were recorded in both D-6 acetone and D-4 methanol. In methanol four resonances were readily discernible for PDA (Figure 2); these all occurred in the aromatic region of the spectrum. The peak at 147.87 ppm arose from the non-protonated aromatic carbon of PDA. This was readily rationalized by its line width (6) and extremely long spin-lattice relaxation time of > 60 seconds (s) (*vide infra*). The other three resonances occurred at 132.35, 130.97, and 129.74 ppm. From intensity considerations and resonance patterns, we reasoned that the peak furthest downfield must be assigned to the C-4 of PDA, whereas the other two resonances must arise from the two equivalent C-2,4 and C-3,5 atoms, respectively.

In acetone the same resonances occurred at 147.18, 132.65, 131.00, and 129.78 ppm. Some of these were substantially broadened, which may result from solvent-solute interactions. Further studies are needed to delineate this.

PDA and BAL. PDA readily reacts with BAL to form a 5-membered ring adduct as shown below (1).



We attempted to undertake two studies of this complex: one study in methanol to monitor the adduct being formed and another in acetone to monitor the T1's of various carbon atoms of BAL before and after the adduct was formed.

The reaction of PDA with BAL was initially undertaken in D-4 methanol (Figure 3). The formation of the adduct had a pronounced effect on the chemical shifts of the PDA resonances. New resonances appeared at 144.4 (1), 131.64 (1), 130.06 (2), and 129.40 (2) ppm, respectively. The number in parentheses reflects the carbon count inferred from peak intensities. This

indicated that one of the carbons shifted as much as 3.4 ppm due to adduct formation. BAL in D-4 methanol exhibited three resonances at 65.77, 45.54, and 30.00 ppm. Formation of the PDA-BAL adduct had a more pronounced effect on the BAL ^{13}C -resonances than on the ^{13}C -resonances of the phenyl carbons of PDA. The new resonances appeared at 64.68, 59.26, and 43.45 ppm. Clearly, adduct formation had resulted in a shift of as much as 13.4 ppm for two of the carbon atoms.

The second study of the PDA-BAL complex was performed in D-6 acetone. The PDA resonances for the complex occurred at 147.06 (1), 132.22 (1), 130.77 (2), and 129.52 (2) ppm, respectively. It appeared that adduct formation in the D-6 acetone solvent had little effect on ^{13}C chemical shifts of PDA resonances. This is a point that should invite further investigation. BAL in D-6 acetone exhibited two observable resonances at 65.46 (triplet) and 45.56 (doublet) ppm when monitored (proton coupled). Unfortunately, the third resonance lay under one of the solvent's (D-6 acetone) resonances. When the PDA-BAL adduct was monitored (proton coupled), the BAL resonances occurred at 64.25 (triplet), 59.36 (doublet), and 40.65 (triplet) ppm. The resonance that shifted from 45.56 to 59.36 ppm was clearly the C-2 carbon of BAL, and the resonance that hardly shifted (65.46 to 64.25 ppm) was the C-1 carbon. The remaining resonance was attributed to the C-3 of BAL. This interpretation was directly supported by the data in the previous study in which D-4 methanol was the solvent.

In order to ascertain the relative mobility (via the rotational correlation time, τ_R) of certain carbon atoms of BAL in the "free" form and in the adduct, T1 values for BAL and the BAL-PDA adduct were obtained. In the free form, the T1's of C-1 and C-2 of BAL in acetone (336 mM) were 5.6s and 8.1s, respectively. The acetone resonance obscured the ^{13}C resonance of BAL C-3. In the BAL-PDA complex in D-6 acetone (336 mM), T1 values of 4.4, 8.1, and 5.0 s, respectively, were observed for BAL C-1, C-2, and C-3. Thus, in acetone, the T1's of BAL appear to be unaffected by adduct formation.

The T1's of the PDA carbon atoms of the PDA-BAL complex in D-6 acetone were 53 s (129.4 ppm) 2.7 s (132.4 ppm), 6.1 s (130.8 ppm) and 5.6 s (129.5 ppm). These were not compared with the ^{13}C -T1 values of free PDA. It is interesting to note that the T1 of C-1 PDA was approximately 1 minute.

PDA and Glutathione

Glutathione is a tripeptide (γ -L-glutamyl-L-cysteinyl-glycine) that is found at a concentration of 2 mM in the human red blood cell. It contains a cysteine residue that has a sulfhydryl group which can readily react with PDA. Our results given below, along with the peak integration data of the various spectra (not presented here) indicate that glutathione forms a 2:1 adduct with PDA in ethanol. This result is similar to the recently published data concerning a glutathione-vanadium complex (7).

The resonances for glutathione [333 mM in a water/deuterated water (3:1) mixture] at pH 7.2 are listed in Table 1. The resonance assignments are based on literature data for glutathione (7) and related dipeptides composed of Glu (8).

O'Connor et al--8

Table 1. ^{13}C Chemical Shift Data for Glutathione

Shift in ppm		Assignment
177.1		
175.8	Carbonyl	
174.8	Carbon Atoms	Cys
172.5		
56.4		Cys C^α
54.9		Glu C^α
44.1	Aliphatic	Gly C^α
32.1	Carbon Atoms	Glu C^γ
26.9		Glu C^β
26.5		Cys C^β

Table 2 lists chemical shifts which were obtained after glutathione formed a 2/1 adduct with PDA (165 mM) sample, pH 7.4, in a water/deuterated water mixture. External TMS in a capillary was used as a reference. Figure 4 shows the ^{13}C -NMR spectra of glutathione and glutathione reacted with PDA.


Table 2. ^{13}C Chemical Shift Data for PDA-Glutathione Adduct

Shift in ppm		Assignment
176.8	Carbonyl Carbon Atoms	Cys
175.4		
174.6		
172.0		
139.4	PDA Carbon Atoms	
132.4		
130.1		
55.5	Aliphatic Carbon Atoms	Cys C^α
54.8		Glu C^α
44.2		Gly C^α
33.5		Cys C^β
32.2		Glu C^γ
26.9		Glu C^β

It can readily be seen that the carbon atoms whose chemical shifts were perturbed upon formation of the adduct are those of cysteine. These results indicate that the sulfhydryl groups protrude, unhindered, into the aqueous media and that the adduct formation does little to disturb the structure of the tripeptide. We also noted in our studies that when DMSA was added to the solution, the PDA-glutathione complex was readily abolished.

PDA-DMSA

DMSA is a compound that contains vicinal sulfhydryl groups.



These groups readily react with PDA to form a thermodynamically stable 5-membered ring. DMSA is only sparingly soluble in D-6 acetone. However, it reacted with PDA in acetone to form the 1:1 DMSA-PDA adduct. A signal for the methine carbons of DMSA in the complex was observed at 59.3 ppm. Another signal also appeared at 57.5 ppm, and it appeared to increase substantially with time. It probably was an oxidation product of DMSA.

CONCLUSIONS

We have shown that when PDA is added to red blood cells, it will penetrate the RBC and bind (react) with some component in the cell interior. Our NMR results indicate that the intracellular component is probably glutathione.

RECOMMENDATIONS

A number of NMR experiments need to be performed to characterize further the PDA-adducts (BAL, DMSA). Examples are:

- (1) Effect of solvent on the chemical shifts of these complexes. How important is a protic or aprotic solvent?
- (2) T1's should be determined for the PDA-BAL complex in solvents other than acetone. Again, a protic solvent may have a different effect.
- (3) ^{75}As -NMR of these species.
- (4) Undertake 2D-NMR studies of these various complexes.

REFERENCES

1. Webb JL. Arsenicals. In: Enzyme and metabolic inhibitors. New York: Academic Press, 1966; Volume 3.
2. Gates M, Williams JW, Zapp JA. Arsenicals. In: Chemical warfare agents and related chemical problems. Summary Technical Report of Division 9, National Research Committee, Office of Scientific Research and Development, 1946; Volume 1.
3. Prentiss AM. Chemicals in war. New York: McGraw-Hill Book Company, 1937; 177-200.
4. Board PG, Agar NS. Glutathione metabolism. In: Red blood cells of domestic mammals. NS Agar and PG Board, eds. New York: Elsevier, 1983, pp 253-270.
5. Jocelyn PC. Biochemistry of the SH-group. New York, Academic Press, 1972, pp 240-243.
6. Oldfield E, Norton RS, and Allerhand A. Studies of individual carbon sites of proteins in solution by natural abundance carbon-13 nuclear magnetic resonance spectroscopy. J Biol Chem, 1975; 250, 6368-6380.
7. Delfini M, Gaggelli E, Lepsi A, and Valensi G. Nuclear magnetic resonance study of the oxovanadium (iv)-(Glutathione)₂ complex. Inorg Chim Acta, 1985; 107, 87-89.
8. Lianon HK, Dillek, (and Pavia AA. Int. J. Peptide Protein Res, 1986 (in press).

LIST OF FIGURES

Figure		Page
1	Scatchard Plot of PDA Binding to Hemoglobin	15
2	Proton-Decoupled ^{13}C -NMR Spectrum of the Aromatic Region of PDA in D-6 Acetone (upper spectrum) and D-4 Methanol (lower spectrum)	16
3	Proton-Decoupled ^{13}C -NMR Spectrum of the Aromatic Region of Free PDA (upper spectrum) and PDA Reacted with BAL (lower spectrum)	17
4	Proton-Decoupled ^{13}C -NMR Spectrum of Native Glutathione (upper spectrum) and Glutathione Reacted with PDA (lower spectrum)	18




Figure 1. Scatchard Plot of PDA Binding to Hemoglobin

O'Connor et al--16




Figure 2. Proton-Decoupled ^{13}C -NMR Spectrum of the Aromatic Region of PDA in D-6 Acetone (upper spectrum) and in D-4 Methanol (lower spectrum).




Figure 3. Proton-Decoupled ^{13}C -NMR Spectrum of the Aromatic Region of Free PDA (upper spectrum) and PDA reacted with BAL (lower spectrum)




Figure 4. Proton-Decoupled ^{13}C -NMR Spectrum of Native Glutathione (upper spectrum) and Glutathione Reacted with PDA (lower spectrum)

O'Connor et al--19
OFFICIAL DISTRIBUTION LIST

Commander
US Army Medical Research
and Development Command
ATTN: SGRD-RMS/Mrs. Madigan
Fort Detrick, MD 21701-5012

Defense Technical Information Center
ATTN: DTIC/DDAB (2 copies)
Cameron Station
Alexandria, VA 22304-6145

Office of Under Secretary of Defense
Research and Engineering
ATTN: R&AT (E&LS), Room 3D129
The Pentagon
Washington, DC 20301-3080

The Surgeon General
ATTN: DASG-TLO
Washington, DC 20310

HQ DA (DASG-ZXA)
WASH DC 20310-2300

Commandant
Academy of Health Sciences
US Army
ATTN: HSHA-CDM
Fort Sam Houston, TX 78234-6100

Uniformed Services University
of Health Sciences
Office of Grants Management
4301 Jones Bridge Road
Bethesda, MD 20814-4799

US Army Research Office
ATTN: Chemical and Biological
Sciences Division
PO Box 12211
Research Triangle Park, NC 27709-2211

Director
ATTN: SGRD-UWZ-L
Walter Reed Army Institute
of Research
Washington, DC 20307-5100

Commander
US Army Medical Research Institute
of Infectious Diseases
ATTN: SGRD-ULZ-A
Fort Detrick, MD 21701-5011

Commander
US Army Medical Bioengineering
Research & Development Laboratory
ATTN: SGRD-UBG-M
Fort Detrick, Bldg 568
Frederick, MD 21701-5010

Commander
US Army Medical Bioengineering
Research & Development Laboratory
ATTN: Library
Fort Detrick, Bldg 568
Frederick, MD 21701-5010

Commander
US Army Research Institute
of Environmental Medicine
ATTN: SGRD-UE-RSA
Kansas Street
Natick, MA 01760-5007

Commander
US Army Institute of Surgical Research
Fort Sam Houston, TX 78234-6200

Commander
US Army Research Institute
of Chemical Defense
ATTN: SGRD-UV-AJ
Aberdeen Proving Ground, MD 21010-5425

Commander
US Army Aeromedical Research Laboratory
Fort Rucker, AL 36362-5000

AIR FORCE Office of Scientific
Research (NL)
Building 410, Room A217
Bolling Air Force Base, DC 20332-6448

Commander
USAFSAM/TSZ
Brooks Air Force Base, TX 78235-5000

Head, Biological Sciences Division
OFFICE OF NAVAL RESEARCH
800 North Quincy Street
Arlington, VA 22217-5000